Stroke Discussion Guide

What can I do to reduce my risk of another stroke?

Recurrent stroke is more common than you may think

1 in 10 survivors will **have another stroke** within a year

1 in 5 survivors will **have another stroke** within five years

After a stroke, it's important to be able to speak confidently with your healthcare provider about your post-stroke care. Use this guide to help shape the conversation during your next visit.

How high is my risk of having another stroke?

What are my personal risk factors for another stroke?

What medication(s) am I on to reduce the risk of another stroke? How should I take each medication?

What other treatment options would you consider for my post-stroke care?

WHAT IS SUNHARBO®?

SUNHARBO is a prescription medicine used to lower the risk of stroke in adults with recent:

- stroke caused by a clot that blocks blood flow in the brain (ischemic stroke), **or**
- high-risk mini stroke (transient ischemic attack or TIA).

SUNHARBO is not for strokes caused by clots that come from the heart.

SUNHARBO should be used in combination with another medicine that helps prevent blood clots (antiplatelet medicine).

It is not known if SUNHARBO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Before taking SUNHARBO, tell your healthcare provider about all your medical conditions, including if you:

- plan to have surgery or a medical or dental procedure. You should stop taking SUNHARBO for at least 24 hours before any planned surgery or procedures.
- have liver problems.
- are pregnant or plan to become pregnant. It is not known if SUNHARBO will harm your unborn baby.

Please see additional Important Safety Information on the following page.

Sunharbo® (asundexian) 50 mg tablets

What changes to my daily habits would best support my recovery?

Which healthcare providers should be involved in my ongoing care, and how will we coordinate my plan?

What changes in movement, speech, memory, mood, vision, swallowing, or fatigue should I report?

What activities are safe for me now, including exercise, driving, work, and travel?

Other things you'd like to discuss (Ex. Recovery, Rehabilitation & Planning):

IMPORTANT SAFETY INFORMATION

Females who are able to become pregnant:

- Talk with your healthcare provider about pregnancy planning during treatment with SUNHARBO.
- If you are exposed to SUNHARBO during pregnancy, talk to your healthcare provider and contact Bayer Healthcare Pharmaceuticals Inc. at 1-888-842-2937.

- are breastfeeding or plan to breastfeed. It is not known if SUNHARBO passes into your breast milk. **Do not** breastfeed during treatment with SUNHARBO and for 5 days after your last dose. Talk to your healthcare provider about the best way to feed your baby during treatment with SUNHARBO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. SUNHARBO may affect the way other medicines work, and other medicines may affect how SUNHARBO works.

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

What are the possible side effects of SUNHARBO?

The most common side effect of SUNHARBO is diarrhea. This is not the only possible side effect of SUNHARBO. Call your doctor for medical advice about side effects.

You are encouraged to report side effects or quality complaints of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. For Bayer products, you can report these directly to Bayer by clicking here.

Please see the full Prescribing Information.

© 2026 Bayer
Bayer, the Bayer Cross, and SUNHARBO are registered trademarks of Bayer.
100 Bayer Boulevard Whippany, NJ 07981
PP-Asun-US-0268-1 11/2026

Sunharbo® (asundexian) 50 mg tablets